

AD-A069 567

FLORIDA STATE UNIV TALLAHASSEE DEPT OF STATISTICS

F/G 12/1

TESTS FOR DEPENDENCE.(U)

FEB 79 M HOLLANDER

AFOSR-78-3678

UNCLASSIFIED

FSU-STATISTICSS-M488

AFOSR-TR-79-0543

NL

| OF |

AD
A069567

12
12



END
DATE
FILMED
7-79

DDC

AFOSR-TR. 79-0543

② LEVEL II


TESTS FOR DEPENDENCE

by

Myles Hollander
Florida State University

FSU Statistics Report M488
AFOSR Technical Report No. 78-2

February, 1979
The Florida State University
Department of Statistics
Tallahassee, Florida 32306



Research sponsored by the Air Force Office of Scientific Research, AFSC,
USAF, under Grant AFOSR 78-3678.

AMS 1970 Subject Classifications: Primary 62G10, 62F05.

9 05 18 185

DISTRIBUTION STATEMENT A

Approved for public release;
Distribution Unlimited

AFOSR-TR-79-0243

LEVEL

2

RECEIVED
JUN 1 1980
AFOSR

DEC FILE COPY

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH (AFOSR)
NOTICE OF TRANSMIT TO DEC
This technical report has been reviewed and is
approved for public release IAW AFR 190-12 (7b).
Distribution is unlimited.
A. D. BLOSE
Technical Information Officer

TESTS FOR DEPENDENCE¹

by

Myles Hollander
The Florida State University

Abstract

This paper is prepared as an invited entry for the Encyclopedia of Statistical Sciences, to be edited by Samuel Kotz and Norman L. Johnson and to be published by John Wiley & Sons. It is designed to provide a sound introduction for a reasonably well-informed reader who is, however, not a specialist in tests for dependence. The paper contains references to many tests but emphasizes the parametric test of independence based on Pearson's sample correlation coefficient r and certain nonparametric tests based on ranks. The ranks tests are generally preferable to the test based on r in that they have wider applicability, are much less sensitive to outlying observations, are exact under mild assumptions which do not require an underlying bivariate normal population, and have good efficiency (power) properties.

1. Introduction

Many studies are designed to explore the relationship between two random variables X and Y , say, and specifically to determine whether X and Y are independent or dependent. Some particular examples are:

(i) *Obesity and blood pressure*: Are obesity and blood pressure independent or, for example, do men who are overweight also tend to have high blood pressure? Here X could be the degree of overweight as measured by the ratio of actual body weight to ideal body weight as given in certain standard tables, and Y could be systolic blood pressure.

¹Research sponsored by the Air Force Office of Scientific Research, AFSC, USAF, under Grant AFOSR-78-3678. The United States Government is authorized to reproduce and distribute reprints for governmental purposes.

(ii) *Color and taste of tuna*: Are color and quality of canned tuna independent or perhaps do consumers tend to prefer light tuna? Here X could be a measure of lightness and Y could be a quality score determined by a consumer panel.

(iii) *Infants walking and their IQ*: Is the time until it takes an infant to walk alone independent of the infant's IQ at a later age, or do children who learn to walk early tend to have higher IQs? Here X could be the number of days measured from birth until the infant walks alone, and Y could be the infant's IQ score at age 5.

(iv) *System reliability and the environment*: Is the life length X (say) of a specific system independent of a certain characteristic of the environment, for example, the temperature Y , within which the system operates, or do high temperatures tend to shorten the life length?

One can test the null hypothesis that the two variables X and Y are independent, against alternatives of dependence, using a random sample from the underlying bivariate population. We suppose that such a sample of size n is available, and we denote the sample by $(X_1, Y_1), (X_2, Y_2), \dots, (X_n, Y_n)$. Our assumptions are

A1. The n bivariate observations $(X_1, Y_1), \dots, (X_n, Y_n)$ are mutually independent.

A2. Each (X_i, Y_i) comes from the same bivariate population with continuous distribution function $H(x, y) = P(X \leq x, Y \leq y)$ and continuous marginal distributions $F(x) = P(X \leq x)$ and $G(y) = P(Y \leq y)$.

The hypothesis of independence asserts that

$$H_0: H(x, y) = F(x)G(y), \text{ for all } (x, y), \quad (1)$$

that is, the variables X and Y are independent. Under H_0 , all $2n$ random variables are mutually independent, that is

$$P(X_1 \leq x_1, Y_1 \leq y_1, \dots, X_n \leq x_n, Y_n \leq y_n) = \prod_{i=1}^n F(x_i)G(y_i).$$

ACCESSION for	
NTIS	White Section <input checked="" type="checkbox"/>
DDC	Buff Section <input type="checkbox"/>
UNANNOUNCED	<input type="checkbox"/>
JUSTIFICATION	
BY	
DISTRIBUTION/AVAILABILITY CODES	
Dist.	AVAIL. and/or SPECIAL
A	

When we discuss alternatives to H_0 , we will be assuming that X and Y are dependent so that (1) fails to hold, but we still insist that the independence *between* the n pairs is preserved.

The organization of this paper is as follows. In Section 2 we present the classical test of H_0 based on Pearson's correlation coefficient r . This test assumes, in addition to A_1 and A_2 , that the underlying population is bivariate normal. Section 3 present rank tests of H_0 which do not require the assumption of normality. These rank tests have additional advantages, relative to the test based on r , including wider applicability, relative insensitivity to outlying observations, and desirable efficiency (power) properties. Section 4 illustrates various tests using data relating to color and taste of tuna.

2. A Test Based on Pearson's Correlation Coefficient

The Pearson correlation coefficient r , proposed by the eminent statistician Karl Pearson in 1896, is

$$r = \frac{n \sum_{i=1}^n X_i Y_i - (\sum_{i=1}^n X_i)(\sum_{i=1}^n Y_i)}{([n \sum_{i=1}^n X_i^2 - (\sum_{i=1}^n X_i)^2] [n \sum_{i=1}^n Y_i^2 - (\sum_{i=1}^n Y_i)^2])^{1/2}} \quad (2)$$

The statistic r is the sample correlation coefficient and is an estimator of the corresponding population parameter ρ , the correlation coefficient of the bivariate population defined by $H(x,y)$. Specifically,

$$\rho = \frac{E(XY) - E(X)E(Y)}{\sigma_x \sigma_y} \quad (3)$$

where E denotes expectation, σ_x is the standard deviation of the X population, and σ_y is the standard deviation of the Y population. It can be shown that for all

samples $-1 \leq r \leq 1$, and for all bivariate populations $-1 \leq \rho \leq 1$. When $\rho > 0$, this may be interpreted as X and Y being positively associated (as measured by ρ) and $\rho < 0$ may be interpreted as X and Y being negatively associated (as measured by ρ). Assuming $H(x,y)$ is a bivariate normal cumulative distribution function with correlation ρ , an exact α level test of H_0 versus $\rho \neq 0$ is

$$\begin{aligned} &\text{reject } H_0 \text{ in favor of } \rho \neq 0 \text{ if } |T| \geq t_{\alpha/2, n-2}, \\ &\text{accept } H_0 \text{ if } |T| < t_{\alpha/2, n-2}, \end{aligned} \quad (4)$$

where $t_{\alpha/2, n-2}$ is the upper $\alpha/2$ percentile point of Student's t distribution with $n-2$ degrees of freedom, and

$$T = (n-2)^{1/2} r / (1-r^2)^{1/2}. \quad (5)$$

Since $|T|$ is an increasing function of $|r|$, the test defined by (4) is equivalent to the test which rejects for large values of $|r|$, and the latter is easily derived to be the likelihood ratio test of H_0 versus $\rho \neq 0$ in the model which assumes bivariate normality. (Of course under the bivariate normality assumption, X and Y are independent if and only if $\rho = 0$.)

One-sided tests based on T are readily defined. To test H_0 versus $\rho > 0$, at the α level, reject H_0 if $T \geq t_{\alpha, n-2}$ and accept H_0 if $T < t_{\alpha, n-2}$. To test H_0 versus $\rho < 0$, at the α level, reject H_0 if $T \leq -t_{\alpha, n-2}$ and accept H_0 if $T > -t_{\alpha, n-2}$.

The two-sided test defined by (5), and the corresponding one-sided tests, are exact (i.e., have true Type I error probability equal to the nominal value α) only when the underlying population is bivariate normal. Approximate (for large n) tests of H_0 which do not require the assumption of bivariate normality treat T as a standard normal random variable under H_0 .

For more information on testing independence in this parametric context, see Bickel and Doksum (1977, Section 6.5.A). Devlin, Gnanadesikan, and Kettenring (1975) point out that r is very sensitive to outliers and consider the related problem of robust estimation and outlier detection with correlation coefficients.

In Section 3 we present nonparametric tests of H_0 which are exact without requiring the assumption of bivariate normality.

3. Rank Tests of Independence

Let R_i be the rank of X_i in the joint ranking from least to greatest of X_1, \dots, X_n and let S_i be the rank of Y_i in the (separate) joint ranking from least to greatest of Y_1, \dots, Y_n .

Under assumptions A1 and A2 and H_0 , the vector of X ranks $R = (R_1, \dots, R_n)$ is independent of the vector of Y ranks $S = (S_1, \dots, S_n)$, and both R and S have uniform distributions over the space \mathcal{P} of the $n!$ permutations (i_1, \dots, i_n) of the integers $(1, \dots, n)$. That is, for each permutation (i_1, \dots, i_n) ,

$$P_0\{(R_1, \dots, R_n) = (i_1, \dots, i_n)\} = 1/n!,$$

with the same result holding for (S_1, \dots, S_n) . (The subscript 0 indicates the probability is computed under H_0 .) It follows that rank statistics (i.e., statistics which are solely based on R and S) are distribution-free under H_0 .

One important class of rank statistics for testing H_0 are the *linear* rank statistics of the form

$$L = \sum_{i=1}^n a(R_i)b(S_i) \tag{6}$$

where the "scores" $a(R_i), b(S_i)$ satisfy $a(1) \leq \dots \leq a(n)$, $b(1) \leq \dots \leq b(n)$.

Test based on Spearman's rank correlation coefficient: Making the choice $a(i) = b(i) = i$ in (6), L reduces to

$$M = \sum_{i=1}^n R_i S_i. \quad (7)$$

Then if M is linearly transformed so that the minimum and maximum values are -1 and 1 , we obtain Spearman's rank order correlation coefficient

$$r_s = \frac{12 \sum_{i=1}^n [R_i - (n+1)/2][S_i - (n+1)/2]}{n(n^2 - 1)}. \quad (8)$$

An even simpler formula for computational purposes is

$$r_s = 1 - \frac{6 \sum_{i=1}^n D_i^2}{n^3 - n}, \quad (9)$$

where $D_i = R_i - S_i$. Note also that r_s is obtainable from $r(2)$ by replacing X_i with its X -rank R_i and Y_i with its Y -rank S_i .

The statistic r_s does not estimate ρ as given in (3) but rather it estimates the population parameter

$$\rho_s = 6P\{(X_1 - X_2)(Y_1 - Y_3) > 0\} - 3. \quad (10)$$

It can be shown that for all samples $-1 \leq r_s \leq 1$, and for all bivariate populations $-1 \leq \rho_s \leq 1$. Note that

$$P\{(X_1 - X_2)(Y_1 - Y_3) > 0\} = P(X_1 > X_2, Y_1 > Y_3) + P(X_1 < X_2, Y_1 < Y_3)$$

and when H_0 is true

$$P\{(X_1 - X_2)(Y_1 - Y_3) > 0\} = P(X_1 > X_2)P(Y_1 > Y_3) + P(X_1 < X_2)P(Y_1 < Y_3) = \frac{1}{2} + \frac{1}{2} = \frac{1}{2},$$

so that when H_0 is true, $\rho_s = 0$. In addition, $\rho_s > 0$ may be interpreted as X and Y being positively associated (as measured by ρ_s), and $\rho_s < 0$ may be interpreted as X and Y being negatively associated (as measured by ρ_s). (For further information and interpretation of the parameter ρ_s as a measure of association see Kruskal (1958).)

Under assumptions A1 and A2, an exact α level test of H_0 versus $\rho_s \neq 0$ is

$$\begin{aligned} &\text{reject } H_0 \text{ in favor of } \rho_s \neq 0 \text{ if } |r_s| \geq r_s(\alpha/2, n), \\ &\text{accept } H_0 \text{ if } |r_s| < r_s(\alpha/2, n), \end{aligned} \quad (11)$$

where $r_s(\alpha/2, n)$ is the upper $\alpha/2$ percentile point of the null distribution of r_s . To test H_0 vs. the one-sided alternative $\rho_s > 0$, at the α level, reject H_0 if $r_s \geq r(\alpha, n)$ and accept H_0 otherwise. To test H_0 vs. $\rho < 0$, at the α level, reject H_0 if $r_s \leq -r(\alpha, n)$ and accept H_0 otherwise.

From (9) we see that tests based on r_s are equivalent to tests based on the statistic ED_1^2 . Glasser and Winter give critical values of r_s and ED_1^2 for $n = 4(1)30$. Tables of the complete null distribution of r_s and ED_1^2 are given for $n = 4(1)11$ in Kraft and van Eeden (1968).

Under H_0 , $E(r_s) = 0$, $\text{Var}(r_s) = 1/(n-1)$, and as n gets large, the distribution of $(n-1)^{1/2}r_s$ tends to the standard normal distribution. Thus approximate (for large n) tests of H_0 can be obtained by treating

$$r_s^* = (n-1)^{1/2}r_s \quad (12)$$

as a standard normal variable under H_0 .

Test based on Kendall's rank correlation coefficient: Kendall's rank correlation coefficient can be written as

$$r_k = \frac{2}{n(n-1)} \sum_{i=1}^{n-1} \sum_{j=i+1}^n \xi(X_i, X_j, Y_i, Y_j), \quad (13)$$

where $\xi(a, b, c, d) = 1$ if $(a-b)(c-d) > 0$, and $= -1$ if $(a-b)(c-d) < 0$. When $(X_i - X_j) \cdot (Y_i - Y_j) > 0$ we say the pairs $(X_i, Y_i), (X_j, Y_j)$ are concordant and when $(X_i - X_j)(Y_i - Y_j) < 0$ we say the pairs are discordant. Note that r_k is a rank statistic ($\xi(X_i, X_j, Y_i, Y_j) = \xi(R_i, R_j, S_i, S_j)$ so that one only needs the ranks to compute r_k) but it is not a linear

rank statistic. However, it can be shown (cf. Hájek and Šidák, 1967, Section II.3.1) that, up to a multiplicative constant, Spearman's r_s is the "projection" of Kendall's r_k into the family of linear rank statistics. The statistic r_k estimates the parameter $\tau = 2P\{(X_1 - X_2)(Y_1 - Y_2) > 0\} - 1$. It can be shown that for all samples $-1 \leq r_k \leq 1$, and for all bivariate populations $-1 \leq \tau \leq 1$. When H_0 is true, $\tau = 0$. In addition, $\tau > 0$ may be interpreted as X and Y being positively associated (as measured by τ), and $\tau < 0$ may be interpreted as X and Y being negatively associated (as measured by τ). The reader should note that τ is analogous to the parameter ρ_s (10) estimated by Spearman's r_s . For details of the relationship between ρ_s and τ , see Kruskal (1958).

From (13) we see that tests based on r_k are equivalent to tests based on

$$K = \sum_{i=1}^{n-1} \sum_{j=i+1}^n \xi(X_i, X_j, Y_i, Y_j). \quad (14)$$

Under assumptions A1 and A2, an exact α level test of H_0 versus $\tau \neq 0$ is

$$\begin{aligned} &\text{reject } H_0 \text{ in favor of } \tau \neq 0 \text{ if } |K| \geq k(\alpha/2, n), \\ &\text{accept } H_0 \text{ if } |K| < k(\alpha/2, n), \end{aligned} \quad (15)$$

where $k(\alpha/2, n)$ is the upper $\alpha/2$ percentile point of the null distribution of K .

To test H_0 vs. $\tau > 0$, at the α level, reject H_0 if $K \geq k(\alpha, n)$ and accept H_0 otherwise. To test H_0 vs. $\tau < 0$, at the α level, reject H_0 if $K \leq -k(\alpha, n)$ and accept H_0 otherwise. Kaarsemaker and van Wijngaarden (1953) give tables of the null distribution of K for $n = 4(1)40$. See also Table A.21 of Hollander and Wolfe (1973). Extended tables up to $n = 100$ are made available on request by D.J. Best, see Best (1973).

Under H_0 , $E(K) = 0$, $\text{Var}(K) = n(n-1)(2n+5)/18$, and as n gets large, the standardized distribution of K tends to the standard normal distribution. Thus approximate (for large n) tests of H_0 can be obtained by treating

$$K^* = K/[n(n-1)(2n+5)/18]^{1/2} \quad (16)$$

as a standard normal variable under H_0 .

Ties: Although assumption A2 precludes the possibility of ties, ties may occur in practice. One method of treating ties, when dealing with rank statistics, is to replace R_i by R_i^* (the average of the ranks that X_i is tied for), S_i by S_i^* (the average of the ranks that Y_i is tied for), compute the rank statistic using the R^* 's and S^* 's, and refer it to the appropriate null distribution tables derived under the assumption of continuity. This, however, yields only an approximate, rather than an exact test.

Exact conditional tests, in the presence of ties, can be performed but they are computationally tedious. See, for example, Lehmann (1975, Section 7.3). For more information on ties, see Hájek (1969, Chapter VII).

Advantages of rank tests: Advantages of rank tests, as compared to the parametric test based on r , include:

1. Wider applicability - To compute a rank statistic, we need only know the ranks, rather than the actual observations.
2. Outlier insensitivity - Rank statistics are less sensitive than r to wildly outlying observations.
3. Exactness - Tests based on rank statistics are exact under the mild assumptions A1 and A2, whereas the significance test based on r is exact only when $H(x,y)$ is bivariate normal.
4. Good efficiency properties - Rank tests of H_0 are only slightly less efficient than the normal theory test based on r under the underlying bivariate population is normal (the home court of r), and they can be mildly and wildly more efficient than r when the underlying bivariate population is not normal. Of course, the efficiency question is complicated as it depends both on the specific rank test under

consideration and the specific measure of efficiency used. Roughly speaking, for large n and dependency alternatives "close" to the null hypothesis, the tests based on r_s and r_k sacrifice nine percent of the information in the sample, as compared to the test based on r , under the underlying population in bivariate normal, and can be much more efficient for certain non-normal populations. For more details on efficiency and power, see Lehmann (1975), Section 7.5E) and Hájek and Šidák (1967, Section VII. 2.4), and the references therein.

Other rank tests: A "normal scores" rank test studied by Fieller and Pearson (1961) and Bhuchongkul (1964) is particularly noteworthy. The normal scores test statistic for independence is a linear rank statistic of the form (6) with $a(i) = b(i) = EV_n^{(i)}$ where $V_n^{(1)} < \dots < V_n^{(n)}$ is an ordered sample of n observations from the standard normal distribution. For a suitable choice of the definition of efficiency and a suitable choice of the nature of dependency alternatives, the normal scores test of independence and the test based on r are equally efficient under "normality" and Srivastava (1973) has shown that the normal scores test is more efficient than the test based on r for "all" (i.e. subject to mild regularity) other cases.

References to other nonparametric tests of independence can be found in Sections 8.1 and 10.2 of Hollander and Wolfe (1973) and in Section 7.5D of Lehmann (1975).

3. Example

The following example is based on data of Rasekh, Kramer, and Finch (1970) in a study designed to ascertain the relative importance of the various factors contributing to tuna quality and to find objective methods for determining quality parameters and consumer preference. Table 1 gives values of the Hunter L measure of lightness, along with panel scores for nine lots of canned tuna. The original consumer panel scores of excellent, very good, good, fair, poor, and unacceptable were converted to the numerical values of 6, 5, 4, 3, 2, and 1, respectively. The panel scores in Table 1

are averages of 80 such values. The Y random variable is thus discrete, and hence the continuity portion of assumption A2 is not satisfied. Nevertheless, since each Y is an average of 80 values, we need not be too nervous about this departure from assumption A2.

It is suspected that the Hunter L value is positively associated with the panel score. Thus we will illustrate the one-sided tests of H_0 versus positive association, based on r , r_s , and r_k . The reader will soon see that all three tests reach the same conclusion, i.e., there is positive association between the Hunter L value and the panel score.

Table 1. Hunter L values and consumer panel scores for nine lots of canned tuna.

Lot	Hunter L Value (X)	Panel Score (Y)
1	44.4	2.6
2	45.9	3.1
3	41.9	2.5
4	53.3	5.0
5	44.7	3.6
6	44.1	4.0
7	50.7	5.2
8	45.2	2.8
9	60.1	3.8

Source. J. Rasekh, A. Kramer, and R. Finch (1970).

Test based on r: From Table 1, we easily calculate $\sum X_i Y_i = 1584.88$, $\sum X_i = 430.3$, $\sum Y_i = 32.6$, $(\sum X_i)^2 = 185158.09$, $(\sum Y_i)^2 = 1062.76$, $\sum X_i^2 = 20843.11$, $\sum Y_i^2 = 125.90$, and from (2) and (5) with $n = 9$, we obtain $r = .57$ and $T = 1.84$. Referring $T = 1.84$ to a t-distribution with 7 degrees of freedom yields a one-sided P value of .054. Thus the test based on r leads to the conclusion that

the Hunter L lightness variable and the panel score variable are positively associated.

The large sample approximation refers $T = 1.84$ to the standard normal distribution yielding an approximate P value of .034.

Test based on r_s : We use Table 2 to illustrate the computation of r_s .

Table 2. Computation of r_s for the canned tuna data

Lot	R	S	D	D^2
1	3	2	1	1
2	6	4	2	4
3	1	1	0	0
4	8	8	0	0
5	4	5	-1	1
6	2	7	-5	25
7	7	9	-2	4
8	5	3	2	4
9	9	6	3	9
				$\Sigma D^2 = 48$

From (9) with $n = 9$ we obtain

$$r_s = 1 - \frac{6(48)}{(9)^{3-9}} = .60.$$

Referring $r_s = .60$ to Table J of Kraft and van Eeden (1968) yields a one-sided P value of .048. Thus the test based on r_s leads to the conclusion that the Hunter L lightness variable and the panel score variable are positively associated.

From (12) we see that the large sample approximation refers $r_s^* = (8)^{\frac{1}{2}}(.6) = 1.70$ to the standard normal distribution yielding an approximate P value of .045. This is in good agreement with the exact P value of .048 based on r_s .

Test based on r_k : Table 3 contains the $\xi(X_i, X_j, Y_i, Y_j)$ values used to compute r_k . For example, the $i = 2, j = 5$ entry in Table 3 is a "-1" because $X_2 > X_5$

Table 3. $\xi(X_i, X_j, Y_i, Y_j)$ values for canned tuna data

$j \backslash i$	1	2	3	4	5	6	7	8
2	1							
3	1	1						
4	1	1	1					
5	1	-1	1	1				
6	-1	-1	1	1	-1			
7	1	1	1	-1	1	1		
8	1	1	1	1	-1	-1	1	
9	1	1	1	-1	1	-1	-1	1

and $Y_2 < Y_5$, yielding $(X_2 - X_5)(Y_2 - Y_5) < 0$ and thus $\xi(X_2, X_5, Y_2, Y_5) = -1$. Summing the 1's and -1's of Table 3 yields $K = 16$ and from (13), $r_k = .44$. Referring $K = 16$ to Table A.21 of Hollander and Wolfe (1973) yields a one-sided P value of .060. Thus there is evidence that the Hunter L lightness variable and the panel score variable are positively associated.

To apply the large sample approximation we compute, from (16), $K^* = 1.67$ yielding an approximate P value of .048. This is in good agreement with the exact P value of .060 based on K .

REFERENCES

- Best, D. J. (1973). Extended tables for Kendall's tau. Biometrika 60, 429-430.
- Bhuchongkul, S. (1964). A class of nonparametric tests for independence in bivariate populations. Ann. Math. Statist. 35, 138-149.
- Bickel, P.J. and Doksum, K.A. (1977). Mathematical Statistics: Basic Ideas and Selected Topics. Holden-Day, San Francisco.
- Delvin, S.J., Gnanadesikan, R., and Kettenring, J.R. (1975). Robust estimation and outlier detection with correlation coefficients. Biometrika 62, 531-545.
- Fieller, E.C. and Pearson, E.S. (1961). Tests for rank correlation coefficients II. Biometrika 48, 29-40.
- Glasser, G.J. and Winter, R.F. (1961). Critical values of the coefficient of rank correlation for testing the hypothesis of independence. Biometrika 48, 444-448.
- Hájek, J. (1969). Nonparametric Statistics. Holden-Day, San Francisco.
- Hájek, J. and Šidák, Z. (1967). Theory of Rank Tests. Academic Press, New York.
- Hollander, M. and Wolfe, D. A. (1973). Nonparametric Statistical Methods. Wiley, New York.
- Kaarsemaker, L. and van Wijngaarden, A. (1953). Tables for use in rank correlation. Statist. Neerl. 7, 41-54.
- Kendall, M.G. (1938). A new measure of rank correlation, Biometrika 30, 81-93.
- Kendall, M.G. (1970). Rank Correlation Methods, 4th ed. Griffin, London.
- Kraft, C.H. and van Eeden, C. (1968). A Nonparametric Introduction to Statistics. Macmillan, New York.
- Lehmann, E.L. (1975). Nonparametrics: Statistical Methods Based on Ranks. Holden-Day, San Francisco.
- Rasekh, J., Kramer, A., and Finch, R. (1970). Objective evaluation of canned tuna sensory quality. J. Food Sci. 35, 417-423.
- Spearman, C. (1904). The proof and measurement of association between two things. Amer. J. Psychol. 15, 72-101.
- Srivastava, M.S. (1973). On a class of nonparametric tests for independence - bivariate case. Can. Math. Bull. 16, 337-342.

UNCLASSIFIED
SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE

1. REPORT NUMBER FSU No. M488 AFOSR NO. 78-1	2. GOV'T. ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE TESTS FOR DEPENDENCE	5. TYPE OF REPORT & PERIOD COVERED Technical Report	
	6. PERFORMING ORGAN. REPORT NUMBER FSU Statistics Report M488	
7. AUTHOR(s) Myles Hollander	8. CONTRACT OR GRANT NUMBER (s) AFOSR 78-3678	
9. PERFORMING ORGANIZATION NAME & ADDRESS The Florida State University Department of Statistics Tallahassee, Florida 32306	10. PROGRAM, ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS	
11. CONTROLLING OFFICE NAME AND ADDRESS Air Force Office of Scientific Research Bolling Air Force Base, D.C. 20032	12. REPORT DATE February, 1979	
	13. NUMBER OF PAGES 14	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office).	15. SECURITY CLASS (of this report) Unclassified	
	15a. DECLASSIFICATION/DOWNGRADING SCHEDULE	
16. DISTRIBUTION STATEMENT (of this paper) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the Abstract).		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS Hypothesis tests, independence, dependence, ranks.		
20. ABSTRACT		

This paper is prepared as an invited entry for the Encyclopedia of Statistical Sciences, to be edited by Samuel Kotz and Norman L. Johnson and to be published by John Wiley & Sons. It is designed to provide a sound introduction for a reasonably well-informed reader who is, however, not a specialist in tests for dependence. The paper contains references to many tests but emphasizes the parametric test of independence based on Pearson's sample correlation coefficient r and certain nonparametric tests based on ranks. The ranks tests are generally preferable to the test based on r in that they have wider applicability, are much less sensitive to outlying observations, are exact under mild assumptions which do not require an underlying bivariate normal population, and have good efficiency (power) properties.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

19. REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM	
1. REPORT NUMBER AFOSR-TR-79-0543	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER	
4. TITLE (and Subtitle) TESTS FOR DEPENDENCE	5. TYPE OF REPORT & PERIOD COVERED Interim rept.		
7. AUTHOR(s) Myles/Hollander	6. PERFORMING ORG. REPORT NUMBER FSU-STATISTICS-M788 7. CONTRACT OR GRANT NUMBER(s) TR-78-2		
9. PERFORMING ORGANIZATION NAME AND ADDRESS The Florida State University Department of Statistics Tallahassee, Florida 32306	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 61102F 2304/A5		
11. CONTROLLING OFFICE NAME AND ADDRESS Air Force Office of Scientific Research/NM Bldg. 410 Bolling AFB, DC 20332	12. REPORT DATE February 1979		
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) 12 17p.	13. NUMBER OF PAGES 15		
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		15. SECURITY CLASS. (of this report) UNCLASSIFIED	
15a. DECLASSIFICATION/DOWNGRADING SCHEDULE			
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)			
18. SUPPLEMENTARY NOTES			
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Hypothesis tests, independence, dependence, ranks.			
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) This paper is prepared as an invited entry for the <u>Encyclopedia of Statistical Sciences</u> , to be edited by Samuel Kotz and Norman L. Johnson and to be published by John Wiley & Sons. It is designed to provide a sound introduction for a reasonably well-informed reader who is, however, not a specialist in tests for dependence. The paper contains references to many tests but emphasizes the parametric test of independence based on Pearson's sample correlation coefficient r and certain nonparametric tests based on ranks. The ranks tests are generally preferable to the test based on r in			

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

20. Abstract continued.

that they have wider applicability, are much less sensitive to outlying observations, are exact under mild assumptions which do not require an underlying bivariate normal population, and have good efficiency (power) properties.

UNCLASSIFIED